DOCUMENTATION PROCEDS FOR HAZARD RANKING SYSTEM

INSTRUCTIONS: The purpose of these records is to provide a convenient way to prepare an auditable record of the data and documentation used to apply the Hazard Ranking System to a given site. The source of information should be provided for each entry and should be a bibliographic-type reference that will allow anyone to find the document used for a given data point. Include the location of the document and consider appending a copy of the relevant page(s) for ease in review by any interested party.

FACILITY NAME:	CAHOKIA DEAD CREEK
LOCATION:	SAUGET IL. (St. CLAIR COUNTY)

1 OBSERVED RELEASE

Contaminants detected off site (5 maximum):

PCB's, chloro aniline, dichloro benzere, copper, mu ganese, lead,

Cyclohexane, chlorotherol, aliphatic hydrocarbons, silver, nickel,

cusemic, cadmium, phosphorous

Reasoning by which the presence of the detected contaminants can be attributed to the facility: The last the second contaminants can be

Found in IEPA monitoring wells surrounding the site Wastes corresponds to waste of several inclustries in the area Site is surrounded by various inclustries surce: "A Preliminary Hydrogeologic Investigation in the Northern Portion of Dead Creek & Vicinity" By Ron St. John, 4/81, IEPA Report pp. 34-39

2 ROUTE CHARACTERISTICS

Depth to Aquifer of Concern,

Name/description of aquifers(s) of concern:

NOT APPLICABLE (N/A) - OBSERVED RELEASE (O.R.)

Depth(s) from the ground surface to the highest seasonal level of the saturated zone [water table(s)] of the aquifer of concern:

N/A - O.R.

Depth from the ground surface to the lowest point of waste disposal/ storage:

N/A - O.R.

Net Precipitation

Mean annual or seasonal precipitation:

N/A - O.R.

Mean annual lake or seasonal evaporation: (

N/A - O.R.

Net precipitation (subtract the above figures):

N/A - O.R.

Permeability of Unsaturated Zone

Soil type in unsaturated zone:

N/A - O.R.

Permeability associated with soil type:

N/A - U.R.

Physical State

Physical state of waste at time of disposal (or generated gases):

N/A - O.R.

3 CONTAINMENT

Containment

Method(s) of waste or leachate containment evaluated:

N/A - O.R.

Method with highest score:

N/A - O.R.

4 WASTE CHARACTERISTICS

Toxicity and Persistence

PERSISTENCE Compound(s) evaluated: 3 PCB's Chloro aniline di chloro benzene cyclohexane

Compound with highest score:

Source: SAX & NEPA DATA

S cores

Bazardous Waste Quantity

Total quantity of hazardous waste at the facility (excluding those with a containment score of 0):

UNKNOWN - AREA HAS BEEN USED AS SEVERAL LANDFILLS SINCE A37 AND NO RECORDS ON AMOUNT HAS BEEN KEPT

Basis of estimating and/or computing waste quantity:

Source: "A Preliminary Hydrogeologic Investigation in the Northern Portion of Recid Creek & Vicinity " By Ron St. John, 4/81, DEPA Report

SCORES A (O)

ida Fis

Ground Water Use

Use(s) of aquifer(s) of concern within a 3-mile radius of the facility: ...

Industrial / Commercial - Source: IL. STATE NATER SURVEY Mississippi River is major water source in the area

Distance to Nearest Well

Location of nearest well drawing from aquifer of concern or occupied. · building not served by a public water supply: ______

< 2000 ft : Monsanto well upgradient to GW flow but couses ame of depression in water table - contaminated well</p>

Distance to above well or building: SCORES A (9) 4 2000 ft.

Population Served by Ground Water Wells Within a 3-Mile Radius

Identified public-supply well(s) drawing from aquifer(s) of concern within a 3-mile radius:

NONE Track Supply I'm & used bottled theo

Private - industrial - Amax

Carro Carp

Population served by each above public-supply well and how computed:

75W - 0 Source: ISWS

Computation of land area irrigated by supply well(s) drawing from aquifer(s) of concern within a 3-mile radius, and conversion to population (1.5 people per acre):

(N/A

Total population served by ground water within a 3-mile radius:

Industrial use only Scores A &

recycled pape:

recogn and environment.

1 OBSERVED RELEASE

Contaminants detected in surface water at the facility or downhill from when it (5 maximum):

PCB's, Chloroaniline, dichlorobenzene, cycloherane, Chlorophenol

Reasoning by which the presence of the detected contaminants can be contaminants can be contaminants on be contaminants.

Contaminants deposited directly into waters - Lab. analysis of create downstream & certain company holding pands show waste analysis of the same type
SOURCE: ST. JOHN'S REPORT

2 ROUTE CHARACTERISTICS

Facility Slope and Intervening Terrain

Average slope of facility in percent:

N/A -0: R.

Name/description of nearest downslope surface water:

N/A - O.R.

Average slope of terrain between facility and above-cited surface water body in percent:

N/A - O.R

Is the facility located either totally or partially in surface water?

N/A - OR.

Is the facility completely surrounded by areas of higher elevation?

N/A -OR

1-Year 24-Hour Rainfall in Inches are well more

N/A .- D.R.

Distance to Nearest Downslope Surface Water home warrant

N/A - O.R.

Physical State of Waste

N/A - D.R.

3 CONTAINMENT

Containment

Method(s) of waste or leachate containment evaluated:

N/A - O.R.

Method with highest score:

N/A - O.R.

4 WASTE CHARACTERISTICS

Refer to Ground Water Route

* * *

5 TARGETS

Surface Water Use San Jacks Con-

Use(s) of surface water within 3 miles downstream of the hazardous substance:

Recreation - Children play in create

Scores A 2

Is there tidal influence?

No

Distance to a Sensitive Environment

Distance to 5-acre (minimum) coastal wetland, if 2 miles or less:

> Zmile

Distance to 5-acre (minimum) fresh-water wetland, if I mile or less:

> 1 mile

Distance to critical habitat of an endangered species or national wildlife refuge, if I mile or less:

> Imile

Scares A @

Population Served by Surface Water

Location(s) of public-supply intake(s) within 3 miles (free-flowing bodies) or 1 mile (static water bodies) downstream of the hazardous substance:

None

Population served by each above public-supply intake:

None

Computation of land area irrigated by above-cited intake(s) and conversion to population (1.5 people per acre):

N/A

Name/description of nearest of above water bodies:

Dead Creek

Distance to above-cited intakes, measured in stream miles.

N/A

TOTAL: SCORES A @

1 OBSERVED RELEASE

Contaminants detected:

No specifics - just a survey to get a plus or minus reading (check for spots carbone booksround)

Source: Fit memo 4/14/82 by Don Woods

Date and location of detection of contaminants

3/23/82

North end of Dead Creek - just south of Queeny Ave.

Methods used to detect the contaminants:

Organic Vapor Analysizer

Z-HNU photo-ionizer 10.2 en lamp

11.7 en lamp

Explosimater, Radiation Survey mater. Oz Indicato

Reasoning by which the presence of the detected contaminants can be attributed to the site:

Absorption of contaminants with the soil, a buried pipe was found entering the creek with a tunbid & only liquid coming from it. Midwest Rubber use to pipe their waste into the creek (creek bed is sparsy when you walk on it)

2 WASTE CHARACTERISTICS

Reactivity and Incompatibility

Three most reactive compounds (indicate one used):

Phosphorous

Arsenic

SCORES A D

Three most incompatible pairs of compounds (indicate one used):

Toxicity

Three most toxic compounds (indicate one used):

PCB's

Dichloro phenol

chloroaniline

Hazardous Waste Quantity

Total quantity of hazardous waste:

UNKNOWN

Basis of estimating and/or computing waste quantity:

3 TARGETS

Population Within 4-Mile Radius

Circle radius used, give population, and indicate how determined:

1 to 4 mi / 1/2 to 1 mi

1/4 to 1/2 mi

0 to 1/4 mi

6, 100 people

Screes A (2)

Distance to a Sensitive Environment

Distance to 5-acre (minimum) coastal wetland, if 2 miles or less:

12 Zmites

Distance to 5-acre (minimum) fresh-water wetland, if I mile or less:

> Inile

Distance to critical habitat of an endangered species, if I mile or less:

Land Use

Distance to commercial/industrial area, if I mile or less:

Kymile (22000ff) Topo map & FIT inspection

Scares A 3)

Distance to national or state park, forest, or wildlife reserve, if 2 miles or less:

None

Distance to residential area, if 2 miles or less:

< 1/2 mile -bpo map & FIT inspection Scores A D

Distance to agricultural land in production within past 5 years, if 1 mile or less:

ZY mile topo mop & FIT inspection

SCORES A

D

Distance to prime agricultural land in production within past 5 years, if 2 miles or less:

Z mile topo map & FIT inspection

Scores A D

Is a historic or landmark site (National Register or Historic Places and National Natural Landmarks) within the view of the site?

NO - only the St. Louis Arch

FIRE AND EXPLOSION - PICTURES FROM JEPA FILE OF CREEK BED ON FIRE

CONTAINMENT - WASTE DUMPED WITH NO SEGREGATION SCORES A 3

WASTE CHARACTERISTICS

DRECT EUIDENCE -> no measurements, but photographs of creek bed on fire

SCORES A 3

IGNITI	ABILITY ->	SCORES A D			
	TOXICITY	PERSISTENCE	IGNI TABILITY	REACTIVITY	
Chlorobensene	2	2	3	0 '	
Dichbrophenol	3	4	0	0	
Chloroaniline	3	1 .	1	Ø	
PCB	3	3	0	Q	
Phosphorous	3	1	3	1	
Arsenic	2	1	0	,	
ayclohercane	7	2	3	△	

Source: SAX AND NERA DATA
REACTIVITY - SCORES A D
INCOMPATIBILITY - SCORES A

HAZ. WASTE QUANTITY -> UNKNOWN: SCORES A @

TARGETS

DIST. TO NEADEST POP.

Zoift - 2640ft Scores A 3

Source: U.S.GS. TOPO MAP (DIST N 500 ft)

DIST. TO NEADEST BLOG.

51 - 200 ft Sources A 2

SOURCE: U.SGS. TOPO MAP (DIST ~ 55ft)

DIST TO SENSITIVE BUVIR. -> SCORES A 1

LAND USE - SCORES A 3 (See air section)

POP WIN 2 MILE RADIUS -> 12,239: 16 of E St Louis \$ 1520 blogs (E St Louis: 51700 pop) countral off of map x 3.6 put/buy. \$700/8=6463 people SCORES A (5)

BLOGS. WIN 2 MILRAD. - 1520 counted; 6463/3.8 = 1700 TOTAL: 3220

DIRECT CONTACT

OBSERVED INCIDENT -> Yes, resident's dog rolled in ditch and ched of apparent enemical burns in Aug. 1980 SOURCE: ST JOHN'S REPORT

WHATE CHARACTERISTICS; TORICITY - PCB's = 3 SCORES A 3

DIST. TO CRITICAL HABITAT: SCORES A @